

# **THE OVARIAN INSULIN AND INSULIN-LIKE GROWTH FACTOR SYSTEM WITH AN EMPHASIS ON DOMESTIC ANIMALS<sup>1</sup>**

L.J. Spicer<sup>\*,2</sup> and S.E. Echternkamp<sup>\*\*</sup>

<sup>\*</sup>Department of Animal Science, Oklahoma State University  
Stillwater, OK 74078  
and

<sup>\*\*</sup>U.S. Meat Animal Research Center, Agricultural Research Service, U.S.  
Dept. of Agriculture, Clay Center, NE 68933

Received November 15, 1994

## **ABSTRACT**

Insulin and insulin-like growth factors (IGFs) have direct effects on cultured ovarian cells. These effects include stimulation of granulosa cell mitogenesis, granulosa and luteal cell progesterone production, and thecal cell androgen production and appear similar among species. However, species differences exist with regard to insulin and IGF-I effects on granulosa cell estradiol production. In addition to endocrine effects of insulin and IGFs, IGFs are produced by granulosa, thecal, and luteal cells, allowing for an intraovarian autocrine and paracrine system. Granulosa, thecal, and luteal cells contain receptors for insulin and IGFs, and these receptors appear to mediate the effects of insulin and IGFs. Adding to the complexity of the regulatory role of IGFs is the presence of IGF-binding proteins (IGFBPs) within the ovary. These IGFBPs are produced by granulosa, thecal, and luteal cells, and their production is hormonally regulated. Evidence for a coherent mechanism by which insulin, IGFs, and IGFBPs interact and regulate ovarian function in vivo has yet to be found.

## **INTRODUCTION**

Over 20 years after the determination of the primary amino acid sequence of insulin in 1955 (1), a role for insulin as a regulator of ovarian function was suggested (2). It took an additional 10 years to determine the mechanism of insulin action on ovarian cell steroidogenesis as well as to determine the presence of insulin receptors in ovarian cells (for a review, see Ref. 3). In general, insulin's effects on ovarian cells are positive, stimulating granulosa cell proliferation and the production of progesterone and enhancing luteal cell steroidogenesis regardless of species. Various aspects of insulin's action on ovarian cells will be detailed below.

Related structurally and functionally to insulin, insulin-like growth factor-I (IGF-I) and IGF-II were purified and sequenced in 1978 (4,5). Approximately 45% of the amino acids comprising IGF-I, IGF-II, and insulin are identical; structural similarities across several species are compared in Table 1. Since that time, the presence and function of IGF-I and -II in many tissues of the body have been identified (for reviews, see Refs. 6–9). Among these is their presence and function in the ovary of many species (for reviews, see Refs. 10–13). For example, in cattle and pigs, IGF-I has been found to stimulate granulosa cell proliferation and mitogenesis (14–16) and enhance follicle-stimulating hormone (FSH)-induced steroidogenesis by granulosa cells (15–17). In addition, the presence of high-affinity, low-capacity binding sites for insulin and IGF-I in granulosa cells of pigs and

TABLE 1. HOMOLGY OF IGFs AND INSULIN AMONG SPECIES.<sup>a</sup>

Bovine Insulin/IGF	Ovine	Porcine	Human	Rat
IGF-I (%)	99	100	100	96
IGF-II (%)	99	97	96	96
Insulin (%)	≥97	≥95	≥93	≥95

<sup>a</sup> Compiled from several sources.

cattle has been documented (16–21). Moreover, the production of IGF-I by bovine and porcine granulosa cells appears to be under hormonal control (15,22,23). More recently, hormonally sensitive IGF-binding proteins (IGFBPs) have been found to be secreted by porcine granulosa cells in vitro (24–26). This review will summarize current evidence regarding the nature of the multiple components of the ovarian insulin and IGF system, emphasizing data generated in domestic animals.

### ROLE OF INSULIN AND IGF IN OVARIAN FUNCTION

**Granulosa cell mitogenesis.** Studies in swine, sheep, and cattle have shown stimulatory effects of insulin and IGF-I on granulosa cell proliferation and(or) DNA synthesis (15,16,27–33) and are summarized in Table 2. The ED<sub>50</sub> for the effects of insulin (100 ng/ml) and IGF-I (30 ng/ml) on mitogenesis indicates that IGF-I is a more potent mitogen than is insulin. In cultured porcine granulosa cells, IGF-I appears to enhance the mitogenic effects of other growth factors (29). In further support of a role for IGF-I in follicular growth, concentrations of IGF-I in the follicular fluid of cattle correlate positively with follicular diameter (see later section on “Ovarian Insulin and IGFs”). Few studies have evaluated the dose-response effect of IGF-II on granulosa cell proliferation, although IGF-II does stimulate DNA synthesis in cultured porcine (16,34) and human (35) granulosa cells. In rats, insulin and IGF-I also appear to stimulate granulosa cell mitosis (36–38) and this mitogenic effect of IGF-I is enhanced by FSH (38). Similarly, recent in vitro studies in cattle suggest that FSH and luteinizing hormone (LH) enhance the mitogenic effect of IGF-I in granulosa cells from small (<5 mm) but not large (<10 mm) follicles (33). However, FSH does not alter the stimulatory effect of IGF-I on the proliferation of ovine granulosa cells in vitro (31). In vivo, pregnant mare serum gonadotropin (PMSG)–stimulated prepubertal gilts treated with insulin have increased numbers

TABLE 2. SUMMARY OF THE MITOGENIC ACTION OF IGF-I IN BOVINE, OVINE, AND PORCINE OVARIAN CELLS.

Authors	Ref. No.	Year	Endpoint	Fold Increase	Cell Type	Species
Savion et al.	27	1981	Cell numbers	3	Granulosa	Bovine
Baranao and Hammond	16	1984	[ <sup>3</sup> H]thymidine <sup>a</sup>	5	Granulosa	Porcine
May et al.	29	1988	[ <sup>3</sup> H]thymidine	3	Granulosa	Porcine
			Cell numbers	3	Granulosa	Porcine
Kamada et al.	34	1992	[ <sup>3</sup> H]thymidine	3	Granulosa	Porcine
May et al.	46	1992	Cell numbers	2	Thecal	Porcine
Monniaux and Pisselet	30	1992	[ <sup>3</sup> H]thymidine	2	Granulosa	Ovine
			Cell numbers	2	Granulosa	Ovine
Zhang and Bagnel	32	1993	[ <sup>3</sup> H]thymidine	3	Granulosa	Porcine
			Cell numbers	5	Granulosa	Porcine
Chakravorty et al.	47	1993	[ <sup>3</sup> H]thymidine	4	Luteal	Bovine
Spicer et al.	15	1993	Cell numbers	2	Granulosa	Bovine
Monniaux et al.	31	1994	Cell numbers	2	Granulosa	Ovine

<sup>a</sup> [<sup>3</sup>H]thymidine incorporation, a measure of DNA synthesis.

of small (but not medium or large) follicles (39), whereas increased numbers of small antral follicles are correlated with increased serum IGF-I in prepubertal gilts injected daily with porcine somatotropin (pST; 40). In prepubertal gilts treated with sustained-release implants of pST, increased numbers of medium follicles are correlated with increased serum IGF-I (41), suggesting that the mode of pST treatment may affect the size category of follicle that responds. In cattle, bovine ST (bST) treatment for two estrous cycles increased serum IGF-I concentrations and numbers of small (2 to 5 mm) follicles twofold without affecting serum LH or FSH or the numbers of large or medium follicles (42). Treatment with bST also increased serum IGF-I and the ovulation rate in superovulated beef heifers (43,44). However, insulin treatment (causing a twofold increase in plasma insulin concentrations) given concomitantly with FSH during the superovulation of cattle had no effect on the numbers of antral follicles of any size category (45). Collectively, in vitro and in vivo data support the notion that both insulin and IGF-I act as promoters of follicular growth, especially in pigs and cattle.

There is a paucity of information regarding the effects of insulin and IGFs on thecal and luteal cell proliferation. In cultured porcine thecal cells, IGF-I appears to enhance the mitogenic effects of other growth factors such as epidermal growth factor (EGF) and platelet-derived growth factor (46), but its effects are less pronounced than those observed with granulosa cells (29). Recently, Chakravorty et al. (47) reported that insulin was over 80-fold less potent than IGF-I in stimulating DNA synthesis ( $ED_{50} = 0.5 \mu\text{g/ml}$  vs.  $6.0 \text{ ng/ml}$ ) in cultured bovine luteal cells and that their effects were not additive, suggesting a common mechanism of action. Clarification of the role of insulin and IGFs on thecal and luteal cell proliferation awaits further study.

**Estradiol production.** The effects of insulin, IGF-I, and IGF-II on estrogen production by granulosa cells seem to depend on the species studied. In rats and primates, studies indicate that insulin, IGF-I, and IGF-II can stimulate granulosa cell estradiol production in vitro ( $ED_{50} = 250$  and  $40 \text{ ng/ml}$  for insulin and IGF-I, respectively), in a rank order potency suggestive of an effect mediated via type I IGF receptors (10–13,34,48–51). However, in cattle, insulin is a more potent and efficacious stimulator of estradiol production than is IGF-I (15,20,52,53). In fact, IGF-I at  $100 \text{ ng/ml}$  either had no effect or inhibited basal and FSH-induced estradiol production by bovine granulosa cells collected from small (i.e., 1 to 5 mm) follicles, whereas IGF-I at  $100 \text{ ng/ml}$  was a weak stimulator of FSH-induced estradiol production by granulosa cells from large (i.e.,  $\geq 8 \text{ mm}$ ) follicles (15,52). However, when granulosa cells were cultured in the presence of 1% fetal calf serum (which would contain insulin, IGF-I, and IGF-II), IGF-I and insulin had no effect on basal estradiol production by cells collected from large (12 to 15 mm) bovine follicles (54). More recently, we have observed that IGF-I and -II can inhibit insulin-stimulated estradiol production by granulosa cells of both small and large bovine follicles (20). We have further evaluated low-dose (i.e.,  $\leq 10 \text{ ng/ml}$ ) effects of insulin on bovine granulosa cell estradiol production and find that physiologic levels of insulin stimulate follicular estradiol production by more than threefold (Figure 1; 20). In support of these findings, we observed that, in vivo, insulin treatment (causing a twofold increase in plasma insulin concentrations) during the superovulation of cattle increased follicular fluid estradiol concentrations by fivefold in large follicles (45). Also in cattle, we have observed significant increases in the follicular fluid concentrations of estradiol without a change in the follicular fluid concentrations of IGF-I (55,56) and, conversely, significant increases in follicular fluid IGF-I without a change in follicular fluid estradiol (57). Likewise, significant decreases in follicular fluid IGF-I occurred without a change in follicular fluid estradiol of beef heifers (58). Furthermore, concentrations of IGF-I and estradiol in the follicular fluid of cattle can be either correlated negatively (55) or positively (57,59),




Figure 1. Dose response of insulin on FSH-induced estradiol production by granulosa cells collected from small (1 to 5 mm) bovine follicles. Granulosa cells were cultured for 2 d in the presence of 10% fetal calf serum, washed, and then treated with serum-free medium containing 50 ng/ml ovine FSH and 1  $\mu$ g/ml testosterone with or without the various doses of insulin for an additional 24 hr. Values are means from three separate experiments. Means with different superscripts differ ( $P < 0.05$ ). Reproduced from Spicer et al. (20) with permission.

suggesting that changes in the intrafollicular concentrations of IGF-I are not consistently regulating estradiol production by bovine follicles. In further support of a negative effect of IGF-I on estradiol production, dairy heifers treated with bST for 21 days had 70% higher plasma IGF-I and 30% lower plasma estradiol concentrations (60). It should be pointed out that, *in vivo*, coincident changes in ovarian IGFBP levels may influence IGF-I action (see "Ovarian IGFBPs" for further discussion) and that changes in thecal androgen production may influence estradiol production by the intact follicle.

In contrast to studies in cattle, estradiol production by porcine granulosa cells is not altered (19) or is actually inhibited (61) by insulin *in vitro*, whereas estradiol production by porcine granulosa cells is stimulated by IGF-I (19,62) and to a lesser extent by IGF-II (34). In further support of a role for IGF-I in stimulating follicular estradiol production in pigs, Samaras et al. (63) reported that levels of follicular wall IGF-I mRNA and follicular fluid estradiol were correlated positively ( $r = 0.69$ ), whereas follicular IGF-II mRNA was not significantly correlated with estradiol. Also consistent with these findings, diabetic gilts had significantly lower IGF-I and estradiol concentrations in follicular fluid (64,65), whereas insulin treatment had no effect on follicular fluid estradiol concentrations in PMSG-treated prepubertal gilts (19). Reasons for these apparent differences between cattle and pigs will require further elucidation.

Collectively, *in vitro* and *in vivo* data in cattle support the notion that insulin is a better stimulator of granulosa cell estradiol production than is IGF-I, whereas in pigs, IGF-I is a better stimulator of granulosa cell estradiol production than is insulin. Moreover, the lower potency of insulin than IGF-I in studies with rat and primate granulosa cells implies that insulin, with blood levels of 0.2 to 10 ng/ml, may not be a physiologically relevant promoter of estradiol production in those species.

**Progesterone production.** Numerous studies in several mammalian species including cattle and pigs have shown that insulin, IGF-I, and IGF-II consistently stimulate proges-

terone production by granulosa cells in a rank order potency of IGF-I > IGF-II > insulin (15–17,30,51,66–71). Table 3 summarizes the in vitro effect of IGF-I on progesterone production by ovarian cells of domestic animals. In general,  $\geq 4$  d of treatment with IGF-I results in a greater-fold increase in progesterone production than <4 d of treatment (Table 3). However, in those studies that did not correct for final cell numbers, some of this increase is likely due to increased cell proliferation (Table 3). Average ED<sub>50</sub> values for IGF-I, IGF-II, and insulin are approximately 10, 20, and 100 ng/ml, respectively. These results indicate, like those for estradiol production by rat and human granulosa cells, that the effect of IGF-II and insulin in all mammals studied is mediated at least in part via the type I IGF receptor and that, under normal conditions, insulin may not be a physiologically relevant promoter of progesterone production. In cattle and sheep, concomitant FSH treatment enhances the stimulatory effect of IGF-I on progesterone production (15,30; Table 3). In sheep, it appears that granulosa cells from large (5 to 7 mm) follicles are more responsive to IGF-I in terms of progesterone production than are cells from small (1 to 3 mm) follicles (30,31). Interestingly, IGF-I has no effect on progesterone production by hen granulosa cells (72). We find that, in vivo, concentrations of follicular fluid IGF-I and progesterone are correlated positively in postpartum anestrus cows (55) but not significantly correlated in proestrus cows and heifers (57,59,73). In cyclic gilts, concentrations of follicular wall IGF-I mRNA and follicular fluid progesterone are correlated positively ( $r = 0.52$ ; 63). In addition, it appears that insulin and IGF-I can stimulate progesterone production by bovine (74,75), porcine (76,77), and rat (78,79) thecal cells. Importantly, IGF-I stimulates the synthesis of cholesterol side-chain cleavage enzyme and its mRNA in porcine granulosa cells (80,81) and rat thecal cells (78,79,82), suggesting an effect of IGF-I on gene expression. In summary, it appears that both insulin and IGF-I can stimulate progesterone production by the granulosa cells of sheep, cattle, and pigs (Table 3). However, the effect of insulin appears to be mediated via the Type I IGF receptor.

TABLE 3. CHRONOLOGIC SUMMARY OF THE STIMULATORY EFFECT OF IGF-I ON PROGESTERONE PRODUCTION (EXPRESSED AS FOLD OF CONTROLS) IN BOVINE, OVINE, AND PORCINE OVARIAN CELLS IN VITRO.

Authors	Ref. No.	Year	Fold of Controls	Co-Trt <sup>a</sup>	Cell Type	Size <sup>b</sup> (mm)	Duration of Trt	Species
Baranao and Hammond	16	1984	>15 <sup>c</sup>	None	Granulosa	1–3	4 d	Porcine
Veldhuis et al.	17	1985	8 <sup>d</sup>	None	Granulosa	1–5	2 d	Porcine
Veldhuis et al.	66	1986	23 <sup>d</sup>	None	Granulosa	1–5	4 d	Porcine
			10 <sup>d</sup>	E2	Granulosa	1–5	4 d	Porcine
Schams et al.	68	1988	>14 <sup>d</sup>	None	Granulosa	8–15	5 d	Bovine
			<1 <sup>d</sup>	None	Granulosa	8–15	2 d	Bovine
Holtorf et al.	69	1989	7 <sup>d</sup>	None	Granulosa	>4	5 d	Bovine
Caubo et al.	76	1989	4 <sup>d</sup>	None	Theca	8–10	3 d	Porcine
			7 <sup>d</sup>	hCG	Theca	8–10	3 d	Porcine
McArdle and Holtorf	83	1989	$\leq 3$ <sup>d</sup>	None	Luteal		3 d	Bovine
McArdle et al.	71	1991	3 <sup>d</sup>	hCG	Granulosa	8–15	4 d	Bovine
			15 <sup>d</sup>	None	Granulosa	8–15	4 d	Bovine
Monniaux and Pisselet	30	1992	15 <sup>c</sup>	None	Granulosa	8–15	4 d	Ovine
			20 <sup>c</sup>	FSH	Granulosa	8–15	4 d	Ovine
Sauerwein et al.	85	1992	<2 <sup>d</sup>	None	Luteal		0.5 hr	Bovine
Spicer et al.	15	1993	3 <sup>c</sup>	None	Granulosa	1–5	2 d	Bovine
			5 <sup>c</sup>	FSH	Granulosa	1–5	2 d	Bovine
Monniaux et al.	31	1994	5 <sup>c</sup>	FSH	Granulosa	5–7	4 d	Ovine
			1.5 <sup>c</sup>	FSH	Granulosa	1–3	4 d	Ovine

<sup>a</sup> Co-Trt = co-treatment with IGF-I; E2 = estradiol.

<sup>b</sup> Size of follicle from which cells were collected.

<sup>c</sup> Progesterone production corrected for cell numbers at termination of treatments.

<sup>d</sup> Progesterone production (usually, in nanograms per milliliter) not corrected for cell numbers after treatments.

In vitro studies have shown that insulin and IGF-I can stimulate progesterone production by cells from the corpus luteum of cattle, rats, and rabbits in a rank order potency indicative of a Type I IGF receptor effect (83–87). In vivo, we find that serum progesterone and IGF-I concentrations are correlated positively in postpartum dairy cattle (88,89), and that cows in negative energy balance have lower serum IGF-I and luteal phase progesterone secretion than do cows in positive energy balance (88). Furthermore, bST treatment that increases serum IGF-I concentrations during a superovulation paradigm significantly increases serum progesterone concentrations (43). However, bST treatment that increases serum IGF-I during a normal estrous cycle has no effect on serum progesterone (42,60), implying that systemic IGF-I may influence luteal function only under certain conditions.

**Androgen production.** Studies indicate that both insulin ( $ED_{50} = 200$  ng/ml) and IGF-I ( $ED_{50} = 10$  ng/ml) have stimulatory effects on LH-induced thecal cell androgen biosynthesis in pigs (76,77), rats (78,79,90–92), and humans (93). The mechanism of action of IGF-I on LH-stimulated androgen and progesterone production by rat thecal cells appears to be via the stimulation of thecal cell  $3\beta$ -hydroxysteroid dehydrogenase mRNA (78), cholesterol side chain cleavage mRNA, and  $17\alpha$ -hydroxylase mRNA (79,82). A study with thecal cells isolated from 12 to 15-mm bovine follicles indicated that 2 g/ml of insulin or 100 ng/ml of IGF-I had no effect on basal androstenedione production (54). Similarly, we find that insulin and IGF-I, at concentrations found in the blood of cattle, have no effect on basal androstenedione production but both enhanced LH-induced androstenedione production by two- to sixfold in thecal cells from  $\geq 8$ -mm bovine follicles (75; Figure 2). However, further research is needed to determine the mechanism of action of IGF-I and insulin in bovine thecal cells. In summary, it appears that, in all species studied to date, insulin and IGF-I synergize with LH to promote thecal cell androgen production and that bovine thecal cells appear much more sensitive to insulin than do porcine or rat thecal cells.

**Ovarian insulin and IGFs.** The presence of insulin and IGFs in the ovary was first reported by Hammond et al. (94,95) in the pig. Since that time, the presence of IGFs in the follicular fluid of several other species has been reported, including cattle, sheep, horses, and humans (40,41,55,56,58,73,96–98). In general, concentrations of insulin (0.5 to 10 ng/ml), IGF-I (100 to 500 ng/ml), and IGF-II (20 to 1,000 ng/ml) are equal to or lower in follicular fluid than in plasma (57,96,99,100). However, after a 2-day fast in heifers, follicular fluid concentrations of IGF-I are higher than plasma concentrations (73). Similarly, in fasted humans, follicular fluid concentrations of insulin are higher than plasma concentrations (101). Thus, it appears that nutritional status influences the relationship between ovarian and systemic levels of insulin and IGFs.

In studies with cattle, pigs, and sheep, we and others have found that small ( $\leq 4$  mm) antral follicles have lower concentrations of IGF-I than do large ( $\geq 5$  mm) antral follicles (40,41,57,59,99), and concentrations of IGF-I in follicular fluid are correlated positively ( $r = 0.3$  to  $0.7$ ) with follicular diameter in cattle (55,57,59). Similarly, IGF-I mRNA levels in the follicular walls of cyclic gilts were correlated ( $r = 0.59$ ) with follicular diameter (63). In contrast, we find slightly higher levels of IGF-II in small vs. large bovine (Figure 3) and ovine follicles (98), but no difference between small and medium porcine follicles (40). Whether these differences in concentrations of IGFs between small and large follicles are the result of differences in local biosynthesis or changes in the diffusion of IGFs from plasma remains to be determined. Nonetheless, IGF-I and IGF-II are most likely produced by the ovary. Numerous studies have reported the presence of IGF-I and IGF-II mRNA in ovarian tissue of rats, women, cattle, pigs, and sheep including granulosa, thecal, stromal, and luteal cells (15,50,98,100,102–108). Rat gran-




Figure 2. Effects of insulin and IGF-I on LH-induced androstenedione production by thecal cells collected from large ( $\geq 8$  mm) bovine follicles. Thecal cells were cultured for 2 d in the presence of 10% fetal calf serum, washed, and then treated with serum-free medium containing 100 ng/ml bovine LH with or without the various doses of insulin (A) or IGF-I (B) for an additional 48 hr. Values are means from three separate experiments and are expressed as fold of controls, which averaged  $54 \pm 14$  and  $117 \pm 26$  pg/ $10^5$  cells per 24 hr for Panels A and B, respectively. Means with different superscripts differ ( $P < 0.05$ ). Data are modified from Stewart et al. (75).

ulosa cells appear to contain exclusively IGF-I mRNA, whereas rat thecal cells contain exclusively IGF-II mRNA (105,106). We find that both granulosa and thecal cells of bovine follicles contain IGF-I mRNA (15; Figure 4). Recent studies indicate that IGF-II mRNA exists in porcine ovaries (100) as well as in ovine follicular walls (98). In addition, porcine and bovine granulosa cells secrete IGF-I (15,22,23; Table 4), whereas human granulosa cells predominantly secrete IGF-II (109), exclusively localize IGF-II, and contain IGF-II mRNA (50,110,111). In sheep, more follicular walls have detectable IGF-II mRNA than IGF-I mRNA, as assessed by reverse transcriptase–polymerase chain reaction (98). Also, follicular fluid IGF-II concentrations vary with genotype, whereas IGF-I does not (98).

Some studies have indicated that ST and gonadotropins can stimulate IGF-I and(or) IGF-II production in porcine (22,23; Table 4) and human (109) granulosa cells, respectively. Also, the *in vitro* perfusion of intact rabbit ovaries with bST ( $\geq 10$  ng/ml) for  $>4$  hr stimulated IGF-I secretion coincident with an increase in follicular diameter (112). In contrast, the production of IGF-I by bovine granulosa cells *in vitro* was unaltered by treatment with ST and FSH but was inhibited by treatment with insulin (15; Table 4). Moreover, we are unable to detect IGF-II production by cultured bovine granulosa cells (Spicer and Vernon, unpublished observations). *In vivo*, 20- to 40-d treatment of pST in prepubertal gilts increased serum and follicular fluid IGF-I concentrations but de-




Figure 3. Concentrations of IGF-II in follicular fluid (FFL) of small (1 to 5.9 mm) or large ( $\geq 6.0$  mm) follicles collected from heifers during the preovulatory period. Values were determined by a radioimmunoassay as described in Spicer et al. (98). Asterisk indicates mean differs from 1 to 5.9 mm mean ( $P < 0.05$ ).

creased or had no effect on serum and(or) follicular fluid IGF-II concentrations (100). The treatment of prepubertal gilts with PMSG for 3 d has been shown to either increase (99) or not affect (100) follicular fluid IGF-I concentrations. The relative contribution of changing systemic IGF levels to those in follicular fluid in these and other studies remains to be elucidated. Whether species and(or) methodologic differences exist in terms of hormone-regulated IGF production by granulosa cells also will require further study.

In summary, it appears that, regardless of species, ovarian IGF-I production and IGF-II production are under different control mechanisms and that concentrations of IGF-I increase as follicles develop. However, species differences may exist in terms of the specific cell layer(s) within the follicle that may produce IGF-I and IGF-II and in terms of the hormones that regulate their production.

**Ovarian insulin and IGF receptors.** Specific receptors for insulin and IGFs and their mRNA exist in various types of ovarian cells including granulosa cells (16,18,20,21, 51,71,110,111,113), thecal cells (90,91,93,110,111,114), luteal cells (84,85,87,114-116), and ovarian stromal cells (117-119). Affinity constants for the high-affinity binding sites of insulin, IGF-I, and IGF-II in ovarian tissues range from 1 to 5 nM (Table 5).

Whether numbers of insulin and IGF receptors change as follicles develop is unclear. In swine, granulosa cells from small follicles have fewer insulin receptors than do cells from medium or large follicles (18). Similarly, in women, granulosa and thecal cells from dominant follicles localize more insulin receptor mRNA than do cells from small antral follicles (110,111). However, numbers of total ovarian insulin receptors do not change during the menstrual cycle in women (117). In comparison to insulin receptors, the number of granulosa cell IGF-I receptors is similar among small, medium, and large porcine follicles (19,120). In contrast, we have observed that granulosa cells from large bovine follicles, cultured in vitro, have a 15-fold higher number of IGF-I receptors than






Figure 4. A representative northern blot of ovarian IGF-I mRNA in granulosa, thecal, and luteal tissue of cattle and(or) sheep. Granulosa, thecal, and luteal RNA was subjected to electrophoresis, transferred to nitrocellulose filters, and hybridized with <sup>32</sup>P-labeled IGF-I cDNA as previously described by Spicer et al. (15), except that a bovine cDNA (182) was used instead of a human cDNA. Bli = bovine liver; Bgc = bovine granulosa cells; Btc = bovine thecal cells; Ocl = ovine corpora lutea (Day 11 of estrous cycle); Kb = kilobases.

do cells from small follicles (21). A recent report using the cellular localization of [<sup>125</sup>I]IGF-I within ovaries from bovine fetuses and neonatal calves lends support to our studies in that specific IGF-I binding increased as follicles enlarged from preantral to antral size, concomitant with increases in FSH and LH binding (121). Atretic follicles from these ovaries contained significantly lower numbers of specific IGF-I-binding sites than did nonatretic follicles (121). A similar study using in situ hybridization histochemistry in rat ovaries indicated that IGF-I mRNA is lost whereas IGF-I receptor mRNA is increased as the granulosa cells differentiate and luteinize, whereas IGF-I receptor mRNA disappeared from the granulosa cells of atretic follicles (122). These results are in contrast to the aforementioned studies in pigs in which the number of IGF-I receptors in granulosa cells did not differ between small, medium, and large follicles (19,120) when freshly isolated granulosa cells were used. Such cell populations from freshly isolated follicles are a composite of both live and dead cells from both healthy and atretic follicles (123,124).

TABLE 4. THE OVARIAN GRANULOSA CELL IGF SYSTEM: REGULATION BY LOCAL AND SYSTEMIC FACTORS IN SWINE AND CATTLE.

IGF Component	Animal	Regulation <sup>a</sup>		
		Stimulators	No Effect	Inhibitors
GC IGF-I production <sup>b</sup>	Swine	LH, ST, FSH EGF/TGF $\alpha$ , PDGF	PRL, bFGF	TGF $\beta$
	Cattle	? <sup>c</sup>	ST, FSH	Insulin
GC IGFBP-2 production	Swine	IGF-I, Insulin	IGF-II, EGF, E2	TGF $\beta$ , FSH
	Cattle	?	?	?
GC IGFBP-3 production	Swine	IGF-I, Insulin, PGF $_{2\alpha}$	IGF-II, EGF, E2	TGF $\beta$ , FSH, PGE $_2$
	Cattle	?	?	?
GC IGFBP-4 mRNA	Swine	?	FSH, IGF-I	?
	Cattle	?	?	?
GC IGFBP-5 mRNA	Swine	IGF-I	FSH	?
	Cattle	?	?	?
GC IGF-I receptors	Swine	E2, corticoids	?	?
	Cattle	E2, EGF, FSH	LH, PROG	bFGF
GC IGFBP protease	Swine	FSH	Insulin, PGE $_2$ , ST PRL, E2, PROG, TGF $\beta$	IGF-I, IGF-II
	Cattle	?	?	?

<sup>a</sup> Abbreviations not defined in text: PRL = prolactin; TGF = transforming growth factor; E2 = estradiol; PROG = progesterone; PDGF = platelet-derived growth factor.

<sup>b</sup> GC = granulosa cell.

<sup>c</sup> ? = not yet investigated.

Thus, the presence of dead cells would likely mask any real "size" effect on the number of IGF-I receptors. In further support of the notion that the number of IGF-I receptors in granulosa cells is higher in large vs. small follicles, others (30,48) have shown that steroid production by granulosa cells from large follicles is dramatically increased by IGF-I whereas IGF-I has little or no effect on steroid production by granulosa cells from small follicles. In ovine follicles, histochemically localized [ $^{125}$ I]IGF-I did not differ between small (<2 mm) and large (>2 mm) follicles (125), but this may have been due to a concomitant decrease in IGFBPs (see next section). Recent studies using the cellular localization of IGF-I receptor mRNA in human ovaries indicate that IGF-I receptors are present in granulosa cells of primary and antral follicles of infant ovaries and graafian follicles of adult ovaries (110,111,119,126). Interestingly, human oocytes intensively localize IGF-I receptor (127) and its mRNA (126) whereas rat oocytes localize very little IGF-I receptor mRNA (122). Collectively, ours and other recent studies provide evidence that IGF-I receptors are present in granulosa cells throughout ovarian follicular development, increase as small antral follicles develop into graafian follicles (i.e., as granulosa cells differentiate), and decrease during atresia.

The hormonal regulation of granulosa cell IGF-I receptors is not completely understood. We have observed that numbers of IGF-I receptors in granulosa cells of small

TABLE 5. AFFINITY CONSTANTS AND CROSS-REACTIVITY OF IGF-I, IGF-II, AND INSULIN RECEPTORS IN OVARIAN TISSUE.<sup>a</sup>

Receptor Type	Kd <sup>b</sup> (nM)	Cross-reactivity (%)		
		IGF-I	IGF-II	Insulin
IGF-I	1-3	100	10	0.3
IGF-II	1-5	0.5	100	0.0
Insulin	1-2	1	4	100

<sup>a</sup> Compiled from numerous studies using ovarian tissues.

<sup>b</sup> Dissociation constant of the hormone-receptor interaction.

bovine follicles are increased by EGF, estradiol, and FSH, decreased by basic fibroblast growth factor (bFGF), and unaffected by LH and progesterone (21; Table 4). In contrast, numbers of IGF-I receptors in granulosa cells of large bovine follicles were unaffected by EGF, estradiol, FSH, bFGF, or LH (21). As summarized in Table 4, the number of granulosa cell IGF-I receptors is increased by estradiol (66) and dexamethasone (128) in porcine granulosa cells and by FSH, LH, and  $\beta$ -adrenergic agonists in rat granulosa cells (129,130). The stimulatory effect of EGF, estradiol, and FSH on IGF-I receptors in granulosa cells of small follicles may explain why these hormones synergize with IGF-I to enhance granulosa cell mitogenesis and(or) differentiation. In contrast to IGF-I receptors, IGF-II receptor mRNA and protein are localized within both granulosa and thecal cells of human antral follicles (110,111). In sheep, IGF-II receptors appear to be primarily present in granulosa cells of atretic follicles and in thecal cells of healthy follicles, although specific [ $^{125}$ I] IGF-II binding was also detected in oocytes, stroma, and perivascular cells (131). Determination of whether oocytes of domestic animals contain IGF-I receptors or whether follicles of pigs and cattle contain IGF-II receptors awaits further study.

In rats, large luteal cells have more IGF-I-binding sites than do small luteal cells, and luteal IGF-I receptor mRNA decreases between Days 7 and 15 of pregnancy (87). Whether these changes in luteal IGF-I receptors occur during pregnancy in domestic animals remains to be determined. However, numbers of luteal IGF-I, IGF-II, and insulin receptors do not differ between Days 5 to 7, 8 to 12, or 15 to 18 of the estrous cycle in cattle (85). Insulin receptors are also present in rat luteal cells (84). Additional studies are needed to determine if insulin receptor mRNA or numbers of insulin receptors change during the luteal development of other domestic animals.

In summary, most cell types within the ovary contain insulin and IGF receptors, regardless of species. Limited studies indicate that numbers of IGF-I receptors in granulosa cells are hormonally regulated, increase as the follicle develops, and decrease during atresia. However, the hormonal control of IGF-I receptors in thecal and luteal cells awaits further study. Determination of the regulation of ovarian cell insulin receptor and its mRNA also awaits study.

**Ovarian IGFBPs.** The IGFBPs are high-affinity carrier proteins that function as transport proteins for the IGFs in the systemic circulation. The IGFBPs prolong the half-life of IGFs as well as serve to partition IGFs in extracellular fluids (for reviews, see Refs. 132–134). To date, six unique species of IGFBPs have been identified: IGFBP-1, -2, -3, -4, -5, and -6 (Table 6). The reader is referred to the aforementioned reviews for further details of the chemistry of these IGFBPs. In general, there is about a 50% amino acid homology among the IGFBPs within species and about an 80% amino acid homology for each IGFBP across species. The liver produces IGFBPs along with numerous other tissues including the ovary. Since 1989, when the structure of an IGFBP and its mRNA in ovarian tissue were first reported (135–137), research has implicated the IGFBPs as potential regulators of follicular development. In particular, the IGFBPs in the ovary appear to play an important role in regulating the biologic activity of IGF-I (12,13,137–139). For example, IGFBP-2, -3, -4, and -5 inhibit FSH-induced estradiol synthesis in cultured rat and human granulosa cells by sequestering IGF-I (137,140,141), and the intrabursal administration of IGFBP-3 reduces the number of human chorionic gonadotropin (hCG)-induced ovulations in immature rats (138). Consistent with the idea that IGFBPs inhibit follicular steroidogenesis, levels of follicular fluid progesterone are negatively correlated ( $r = -0.70$ ) with follicular wall IGFBP-2 mRNA (63). Also, follicular fluid estradiol is negatively correlated with follicular fluid IGFBP-2 in ovine follicles ( $r = -0.74$ ; 98) and bovine follicles ( $r = -0.57$ ; 142).

TABLE 6. SUMMARY OF THE SPECIFIC TYPES OF IGFBPs.

IGFBP-1:	25–34 kDa, nonglycosylated (fetal serum, amniotic fluid, endometrium) <sup>a</sup> *Produced by human but not by porcine or rat granulosa cells *Affinity: IGF-I = IGF-II
IGFBP-2:	29–40 kDa, nonglycosylated (fetal serum, CSF <sup>b</sup> ) *Increases with fasting, decreases with ST treatment *Produced by porcine and rat granulosa cells, and by human granulosa and thecal cells *Affinity: IGF-II > IGF-I
IGFBP-3:	150 kDa, glycosylated composed of two subunits, 28 to 53 kDa (adult serum) *ST dependent, decreases with fasting *Produced by luteinized pig granulosa cells, rat corpora lutea, and by human granulosa and thecal cells *Affinity: IGF-I = IGF-II
IGFBP-4:	24–30 kDa (rat serum) *Localized in rat atretic antral and preantral follicles and by human granulosa and thecal cells *Affinity: IGF-I = IGF-II
IGFBP-5:	29–31 kDa (rat serum) *Localized in rat atretic antral and preantral follicles, and corpora lutea, and by human granulosa, thecal and ovarian stromal cells *Affinity: IGF-II > IGF-I
IGFBP-6:	21–32 kDa (adult serum, CSF) *Localized in rat thecal cells; nondetectable in human granulosa and thecal cells *Affinity: IGF-II ≥ IGF-I

<sup>a</sup> Location where IGFBP was first identified.

<sup>b</sup> CSF = cerebrospinal fluid.

Follicular fluid levels of IGFBP-2 as well as other lower molecular weight IGFBPs (but not IGFBP-3), assessed by ligand blotting, decrease as follicles enlarge in sheep (139), pigs (41,143,144), and cattle (142). Similarly, we (98) have observed significantly lower total IGFBP activity and three forms of IGFBPs (IGFBP-2, 27 to 29 and 24 kDa) in large follicles vs. small follicles of ewes 48 hr after prostaglandin  $F_{2\alpha}$  ( $PGF_{2\alpha}$ ) injection and have found that the genotype of ewe had no effect on total IGFBP activity in follicular fluid (as assessed by both exchange assay and ligand blot). The 30% lower total IGFBP activity in large vs. small follicles was not a result of a decrease in IGFBP-3 but rather of a decrease in other lower molecular weight IGFBPs, as previously reported for pigs and cattle (63,142–144). In cyclic gilts, levels of follicular wall IGFBP-2 mRNA and follicular diameter were correlated negatively ( $r = -0.70$ ; 63). Similarly, follicular fluid IGFBP-2 and follicular diameter were correlated negatively in cattle ( $r = -0.81$ ; 142). A decrease in intrafollicular IGFBP activity together with an increase in IGF-I concentrations as follicles develop is thought to provide a coordinated sequence of events that facilitates IGF-I bioavailability for the support of follicular growth and granulosa cell differentiation (12,13). Other data indicate that atresia in ovine follicles <2 mm in diameter is associated with an increase in IGFBP-like activity localized within the granulosa cells and follicular antrum (125). Similarly, atretic follicles in sheep (139), cattle (142), and women (145) contain more IGFBP-2 activity than do healthy follicles. The cause of these changes in follicular fluid IGFBP activity as follicles develop and become atretic is unknown but may involve specific, hormonally regulated proteases that degrade specific IGFBPs (146,147).

Studies evaluating the cellular source of the various ovarian IGFBPs indicate that porcine granulosa, thecal, and(or) luteal cells appear to produce IGFBP-2 and -3 (25,26,147–149). However, in follicular walls (granulosa plus theca) of cyclic gilts, IGFBP-2 mRNA but not IGFBP-3 mRNA could be detected by northern analysis (63). Porcine granulosa cells also appear to produce IGFBP-4 and -5 (150). In comparison, rat granulosa cells appear to produce IGFBP-2, -4, -5, and -6 (151–154). Rat thecal and luteal cells appear to produce only IGFBP-3 (151,155,156). IGFBP-1, produced by the human ovary (13,157), is not found in the porcine or rat ovary. At present, it is uncertain

whether the 29-kDa IGFBP found in the follicular fluid of sheep and cattle is IGFBP-1 or some other IGFBP. Recently, a 31- to 32-kDa IGFBP in bovine follicular fluid has been tentatively identified as IGFBP-5 (158) by western blotting. Verification of these possibilities will require the development of antibodies specific to each of the IGFBPs in each species under investigation. In general, the activity of IGFBP-2 in follicular fluid is greater than or equal to the activity found in plasma, whereas IGFBP-3 activity is lower in follicular fluid than in plasma (98,158–160). How follicular fluid IGFBP-4 and IGFBP-6 activity compares to that found in plasma is unclear because, as mentioned, the specific antibodies necessary to identify them on western blots have yet to be developed for domestic animals. Because of the relatively high IGFBP activity in plasma, it is likely that changing follicular permeability to plasma IGFBPs, in addition to local production, may explain some changes in follicular fluid IGFBP activity as well as alter the bioavailability of intrafollicular IGF-I and -II. Serum IGFBP-2 and IGFBP-3 as well as IGF-I change with feed restriction (161,162), and thus, nutritional state may influence ovarian IGFBP activity. However, like IGF-I and IGF-II, the relative contribution of serum-derived IGFBPs to follicular fluid IGFBP activity remains to be determined.

Additional *in vivo* evidence indicates that total amounts of whole ovarian IGFBP-2 mRNA do not change from Day 11 to 21 of the estrous cycle in gilts, whereas amounts of ovarian IGFBP-3 mRNA decrease during this same interval (149). Interestingly, amounts of IGFBP-2 mRNA are lower whereas amounts of IGFBP-3 mRNA are higher in luteal tissue than in thecal or granulosa cells in gilts (149). Furthermore, amounts of IGFBP-2 increase whereas amounts of IGFBP-3 decrease in the fluid of preovulatory follicles after hCG injection in sows (144). It is speculated by Grimes et al. (148) that the porcine granulosa cells used for their studies luteinize in culture. This may explain why IGFBP-3 is produced by cultured porcine granulosa cells. Thus, together with results from cell culture studies, it appears that IGFBP-2 production decreases and IGFBP-3 production increases as granulosa cells differentiate and luteinize, at least in swine. Additional studies are required to determine if these changes occur in sheep and cattle.

The identification of factors that regulate the production of ovarian IGFBPs has been an active area of research in recent years (Table 4). Of the hormones evaluated, FSH (200 ng/ml) is one of the most potent inhibitors of IGFBP-2 and -3 activity produced by porcine granulosa cells (24). Similarly, in human granulosa cells, IGFBP-2 production is inhibited by hCG and dibutyryl cAMP (163). In cultured rat granulosa cells, total IGFBP activity in spent media was also reduced after FSH ( $\geq 30$  ng/ml) treatment (164,165). The administration of exogenous FSH to cyclic cows decreased the amounts of IGFBP-2 and lower molecular weight IGFBPs in large ( $\geq 8$  mm) estrogen-active follicles but had no effect on these IGFBPs in large estrogen-inactive follicles or on IGFBP-3 in either type of large follicle (142). Prostaglandin  $E_2$  ( $PGE_2$ ) and dibutyryl cAMP have also been shown to inhibit IGFBP activity in cultured porcine (148) and murine granulosa cells (166), suggesting that any cAMP-induced intracellular cascade may inhibit granulosa cell IGFBP activity. Interestingly, the production of IGFBP-1 by human granulosa-luteal cells is stimulated by  $PGE_2$  (157) but inhibited by FSH (167). How gonadotropin-induced cAMP production reduces IGFBP activity is unclear, but recent evidence in rats and pigs suggests that it may involve the induction of specific IGFBP proteases (146,147; Table 4). Further research will be required to determine if other domestic animals have specific ovarian IGFBP proteases.

In contrast to FSH, IGF-I and insulin have been shown to stimulate IGFBP-2 and -3 activity in cultured porcine granulosa cells (25,62) whereas estradiol, EGF, and transforming growth factor- $\beta$  had no significant effect on IGFBP-2 or -3 activity when corrected for changes in cell numbers (24). However, diethylstilbestrol (DES) treatment in

hypophysectomized rats increased ovarian IGFBP-2 mRNA levels yet decreased ovarian IGFBP-6 mRNA levels (152,168). In prepubertal pigs, 20 to 40 d of daily pST injections increased follicular fluid IGFBP-3 activity but did not affect follicular fluid IGFBP-2 activity or whole ovarian IGFBP-2 mRNA, whereas whole ovarian IGFBP-3 mRNA levels decreased significantly (100). These effects of pST were completely eliminated 3 d after gilts were treated with PMSG (100). In contrast, sustained-release implants of pST for 41 d increased IGFBP-2 activity in the follicular fluid of small follicles, had no effect on IGFBP-2 activity in medium follicles, and decreased serum IGFBP-2 activity in prepubertal gilts (41). Thus, the mode of ST treatment may influence the ovarian IGFBP response to ST. Alone, PMSG decreased follicular fluid IGFBP-2 activity but had no effect on the amount of whole ovarian IGFBP-2 or -3 mRNA or on follicular fluid IGFBP-3 activity (100). Why amounts of whole ovarian IGFBP-2 and -3 mRNA do not change in concert with follicular fluid concentrations of IGFBP-2 and -3 is unclear but likely reflects the heterogeneous cell population of whole ovaries as well as the multiple sources of intrafollicular IGFBPs. A comprehensive scheme of how these various hormones and growth factors interplay to regulate ovarian IGFBP activity and gene expression *in vivo* has yet to be devised.

In summary, IGFBP activity in the follicular fluid of pigs, sheep, and cattle decreases as follicles develop and become estrogen active. As follicles undergo atresia and become estrogen inactive, IGFBP activity increases. These changes in IGFBP activity are due to changes in IGFBP-2 and other lower molecular weight IGFBPs and not to changes in IGFBP-3. The hormones that regulate these changes in IGFBP activity are not well characterized for sheep and cattle but, on the basis of studies in rats and pigs, may involve FSH, insulin, and IGF-I (Table 4). The decrease in IGFBP activity together with an increase in IGF-I concentrations as follicles develop likely provides a coordinated sequence of events that facilitates IGF-I and -II bioavailability for the support of follicular growth and increased steroidogenesis by granulosa cells. During luteinization, at least in pigs, IGFBP-3 activity and gene expression increase.

### CONCLUSIONS AND FUTURE DIRECTIONS

Evidence accumulated from *in vitro* studies over the past 10 years clearly indicates that IGF-I at physiologic levels stimulates granulosa cell mitogenesis and steroidogenesis in most species studied. These effects appear to be mediated by specific IGF-I receptors in granulosa cells that increase as follicles develop. Whether IGF-I stimulated mitosis is the reason that follicular diameter and follicular fluid IGF-I are correlated positively *in vivo* remains to be determined. Additional *in vivo* studies have provided evidence that systemic and(or) ovarian IGF-I may be involved in multiple ovulations in cattle (57) but not in sheep (98,169). Comparable data for IGF-II are not available and should be the focus of further research. Strong evidence also exists for a role of insulin and IGF-I in luteal function. Although insulin has ovarian effects that are similar to those of the IGFs, the ED<sub>50</sub> of insulin implies that its effects are not physiologically relevant except in cattle. Recent studies in our laboratory indicate that insulin at physiologic levels indeed may affect ovarian function in cattle.

Both IGF-I and IGF-II are produced by ovarian cells and thus may contribute to an autocrine or paracrine system designed to enhance follicular growth and gonadotropin-induced differentiation. Studies using the *in situ* hybridization of IGF-I and IGF-II mRNAs and insulin, IGF-I, and IGF-II receptor mRNAs will likely reveal the intricacies of the insulin and IGF system that otherwise could not be ascertained. For example, preliminary studies using *in situ* hybridization in ewes indicate that IGF-I mRNA is mainly localized in granulosa, thecal, and luteal cells but the intensity of staining does not

change with follicular or luteal development (170). Elegant studies using coincident *in situ* hybridization of mRNAs and immunocytochemical localization of proteins involved in the IGF system have been conducted with human ovaries (111) and should be a focus of research in domestic animals in the future.

Adding to the complexity of the regulatory role for IGFs is the presence of as many as six specific forms of IGFBPs within the ovary. These IGFBPs are produced by ovarian cells and their production is hormonally regulated. Species differences exist with regard to the specific IGFBP produced and the cell type that produces it, and further research is needed to clarify these differences. A decrease in intrafollicular IGFBP activity together with an increase in IGF-I as follicles develop may facilitate IGF-I bioavailability for the support of follicular growth and for granulosa and thecal cell differentiation. Recent studies indicate that specific proteases may alter porcine (150) and rat (146) ovarian cell IGFBP activity, but more research is required in this area. Future research also should address the role(s) that IGFs and IGFBPs may play in the growth and selection of the follicle(s) destined for ovulation. We have observed that heifers fed to lose body weight had three waves of follicular growth during an estrous cycle whereas heifers fed to gain body weight had primarily two waves of follicular growth (171).

In close anatomical proximity to the ovary is the uterus. Because the uteri of pigs, cattle, and sheep also produce and respond to IGF-I (172–175), future research should determine if communication between the ovarian and uterine IGF system occurs. Current evidence suggests that ovarian and conceptus steroids (i.e., estrogens) regulate the uterine IGF system in rats and pigs (172,176,177). Although additional research will be required to determine if uterine factors (e.g., PGF<sub>2α</sub>, cytokines) regulate the ovarian IGF system, recent data suggest that PGF<sub>2α</sub> stimulates IGFBP-3 production by luteinized porcine granulosa cells (148).

Finally, further research is needed in domestic animals to determine if manipulation of the ovarian IGF system may result in improved reproductive efficiency. As mentioned earlier, Gong et al. (43) found that exogenous bST treatment, which increased serum IGF-I concentrations, increased the number of PMSG-induced ovulations in cattle. In addition, Rieger et al. (178) using cattle and Cognie et al. (179) using sheep have shown that bST injections given concomitantly with FSH injections during a normal superovulatory regimen increased the percentage of transferable embryos. In prepubertal gilts, long-term pST treatment increased the number of antral follicles, which was positively correlated with concentrations of IGF-I in serum (40,41); whether this increased number of follicles will improve reproductive efficiency remains to be determined. It should be emphasized that pST treatment in prepubertal gilts can also inhibit reproductive functions. For example, Bryan et al. (180) observed a significant reduction in the LH responsiveness of porcine granulosa cells cultured *in vitro* from pST-treated gilts. Similarly, we have observed that prepubertal gilts treated with pST have decreased numbers of LH/hCG-binding sites in the granulosa cells of medium follicles (40). In dairy cattle, increased luteal function is associated with increased concentrations of IGF-I in serum during early lactation (88; 181); whether this translates into improved pregnancy rates will require further study.

#### ACKNOWLEDGMENTS/FOOTNOTES

Authors thank Monsanto Co. (St. Louis, MO) for the recombinant bovine IGF-II; F. Gannon (University College Galway, Galway, Ireland) for the bovine cDNA (182); and Paula Cinnamon for secretarial assistance.

<sup>1</sup> Approved for publication by the Director, Oklahoma Agricultural Experiment Station. This research was supported under project H-2088 and by the Cooperative State Research Service, U.S. Department of Agriculture, under agreement No. 93-370203-9023.

<sup>2</sup> To whom reprints should be sent (FAX 405-744-7390).

## REFERENCES

1. Sanger F. La structure de l'insuline. *Bull Ste Chim Biol* 37:33–35, 1955.
2. Channing CP, Tsai V, Sachs D. Role of insulin, thyroxin and cortisol in luteinization of porcine granulosa cells grown in chemically defined media. *Biol Reprod* 15:235–247, 1976.
3. Poretsky L, Kalin MF. The gonadotropic function of insulin. *Endocrine Rev* 8:132–141, 1987.
4. Rinderknecht E, Humble RE. The amino acid sequence of human insulin-like growth factor-I and its structural homology with proinsulin. *J Biol Chem* 253:2769–2776, 1978.
5. Rinderknecht E, Humbel RE. The primary structure of human insulin-like growth factor-II. *FEBS Lett* 89:283–286, 1978.
6. D'Ercole AJ, Underwood LE. Estimation of tissue concentrations of somatomedin C/insulin-like growth factor I. *Methods Enzymol* 146:227–233, 1987.
7. Sara VR, Hall K. Insulin-like growth factors and their binding proteins. *Physiol Rev* 70:591–614, 1990.
8. Humbel RE. Insulin-like growth factors I and II. *Eur J Biochem* 190:445–462, 1990.
9. LeRoith D, Kavanagh VM, Koval AP, Roberts CT Jr. Phylogeny of the insulin-like growth factors (IGFs) and receptors: A molecular approach. *Mol Reprod Devel* 35:332–338, 1993.
10. Adashi EY, Resnick C, Hernandez ER, Svoboda ME, Hoyt E, Clemmons DR, Lund PK, Van Wyk JJ. Rodent studies on the potential relevance of insulin-like growth factor (IGF-I) to ovarian physiology. In: *Growth Factors and the Ovary*, Hirschfield AN (ed.), Plenum Press, New York, pp. 95–105, 1989.
11. Geisthovel F, Moretti-Rojas I, Rojas FJ, Asch RH. Insulin-like growth factors and thecal-granulosa-cell function. *Hum Reprod* 5:785–799, 1990.
12. Hammond JM, Mondschein JS, Samaras SE, Canning SF. The ovarian insulin-like growth factors, a local amplification mechanism for steroidogenesis and hormone action. *J Steroid Biochem Mol Biol* 40:411–418, 1991.
13. Giudice LC. Insulin-like growth factors and ovarian follicular development. *Endocrine Rev* 13:641–669, 1992.
14. Hammond JM, Yoshida K, Veldhuis JD, Rechler MM, Knight AB. Intrafollicular role of somatomedins: Comparison with effects of insulin. In: *Factors Regulating Ovarian Function*, Greenwald GS, Terranova P (eds.), Raven Press, New York, pp. 197–201, 1983.
15. Spicer LJ, Alpizar A, Echterkamp SE. Effects of insulin, insulin-like growth factor I, and gonadotropins on bovine granulosa cell proliferation, progesterone production, estradiol production, and (or) insulin-like growth factor-I production in vitro. *J Anim Sci* 71:1232–1241, 1993.
16. Baranao JLS, Hammond JM. Comparative effects of insulin and insulin-like growth factors on DNA synthesis and differentiation of porcine granulosa cells. *Biochem Biophys Res Commun* 124:484–490, 1984.
17. Veldhuis JD, Furlanetto RW, Junciter D, Garmey J, Veldhuis P. Trophic actions of human somatomedin C/insulin-like growth factor I on ovarian cells: In vitro studies with swine granulosa cells. *Endocrinology* 116:1235–1242, 1985.
18. Otani T, Maruo T, Yukimura N, Mochizuki M. Effect of insulin on porcine granulosa cells: Implications of a possible receptor mediated action. *Acta Endocrinol (Kopenh)* 108:104–110, 1985.
19. Maruo T, Hayashi M, Matsuo H, Ueda Y, Morikawa H, Mochizuki M. Comparison of the facilitative roles of insulin and insulin-like growth factor I in the functional differentiation of granulosa cells: In vitro studies with the porcine model. *Acta Endocrinol (Kopenh)* 117:230–240, 1988.
20. Spicer LJ, Alpizar A, Stewart RE. Evidence for an inhibitory effect of insulin-like growth factor-I and -II on insulin-stimulated steroidogenesis by nontransformed ovarian granulosa cells. *Endocrine* 2:735–739, 1994.
21. Spicer LJ, Alpizar A, Vernon RK. Insulin-like growth factor-I receptors in ovarian granulosa cells: Effect of follicle size and hormones. *Mol Cell Endocrinol* 102:69–76, 1994.
22. Hsu CJ, Hammond JM. Concomitant effects of growth hormone on secretion of insulin-like growth factor I and progesterone by cultured porcine granulosa cells. *Endocrinology* 121:1343–1348, 1987.
23. Hsu CJ, Hammond JM. Gonadotropins and estradiol stimulate immunoreactive insulin-like growth factor-I production by porcine granulosa cells in vitro. *Endocrinology* 120:198–207, 1987.
24. Mondschein JS, Smith SA, Hammond JM. Production of insulin-like growth factor-binding proteins (IGFBPs) by porcine granulosa cells: Identification of IGFBP-2 and -3 and regulation by hormones and growth factors. *Endocrinology* 127:2298–2306, 1990.
25. Grimes RW, Hammond JM. Insulin and insulin-like growth factors (IGFs) stimulate production of IGF-binding proteins by ovarian granulosa cells. *Endocrinology* 131:553–558, 1992.
26. Grimes RA, Samaras SE, Barber JA, Shimasaki S, Ling N, Hammond JM. Gonadotropin and cAMP modulation of IGF binding protein production in ovarian granulosa cells. *Am J Physiol* 262:E495–E503, 1992.



27. Savion N, Liu GM, Laherty R, Gospodarowicz D. Factors controlling proliferation and progesterone production by bovine granulosa cells in serum-free medium. *Endocrinology* 109:409–420, 1981.
28. Dorrington JH, Bendell JJ, Chuma A, Lobb DK. Actions of growth factors in the follicle. *J Steroid Biochem* 27:405–411, 1987.
29. May JV, Frost JP, Schomberg DW. Differential effects of epidermal growth factor, somatomedin-C/insulin-like growth factor I, and transforming growth factor- $\beta$  on porcine granulosa cell deoxyribonucleic acid synthesis and cell proliferation. *Endocrinology* 123:168–179, 1988.
30. Monniaux D, Pisselet C. Control of proliferation and differentiation of ovine granulosa cells by insulin-like growth factor-I and follicle-stimulating hormone in vitro. *Biol Reprod* 46:109–119, 1992.
31. Monniaux D, Pisselet C, Fontaine J. Uncoupling between proliferation and differentiation of ovine granulosa cells in vitro. *J Endocrinol* 142:497–510, 1994.
32. Zhang Q, Bagnell CA. Relaxin stimulation of porcine granulosa cell deoxyribonucleic acid synthesis in vitro: Interactions with insulin and insulin-like growth factor I. *Endocrinology* 132:1643–1650, 1993.
33. Gong JG, McBride D, Bramley TA, Webb R. Effect of recombinant bovine somatotrophin, insulin-like growth factor-I and insulin on the proliferation of bovine granulosa cells in vitro. *J Endocrinol* 139:67–75, 1993.
34. Kamada S, Kubota T, Taguchi M, Wen-Rong H, Sakamoto S, Aso T. Effects of insulin-like growth factor-II on proliferation and differentiation of ovarian granulosa cells. *Horm Res* 37:141–149, 1992.
35. Di Blasio AM, Viganò P, Ferrari A. Insulin-like growth factor-II stimulates human granulosa-luteal cell proliferation in vitro. *Fertil Steril* 61:483–487, 1994.
36. Bendell JJ, Dorrington JH. Epidermal growth factor influences growth and differentiation of rat granulosa cells. *Endocrinology* 127:533–540, 1990.
37. Bley MA, Simon JC, Saraguetta PE, Baranao JL. Hormonal regulation of rat granulosa cell deoxyribonucleic acid synthesis: Effects of estrogens. *Biol Reprod* 44:880–888, 1991.
38. Bley MA, Simon JC, Estevez AG, De Asua LH, Baranao JL. Effect of follicle-stimulating hormone on insulin-like growth factor-I-stimulated granulosa cell deoxyribonucleic acid synthesis. *Endocrinology* 131:1223–1229, 1992.
39. Matamoros IA, Cox NM, Moore AB. Effects of exogenous insulin and body condition on metabolic hormones and gonadotropin-induced follicular development in prepubertal gilts. *J Anim Sci* 69:2081–2091, 1991.
40. Spicer LJ, Klindt J, Buonomo FC, Maurer R, Yen JT, Echternkamp SE. Effect of porcine somatotropin on number of granulosa cell luteinizing hormone/human chorionic gonadotropin receptors, oocyte viability, and concentrations of steroids and insulin-like growth factors I and II in follicular fluid of lean and obese gilts. *J Anim Sci* 70:3149–3157, 1992.
41. Echternkamp SE, Spicer LJ, Klindt J, Vernon RK, Yen JT, Buonomo FC. Administration of porcine somatotropin by a sustained-release implant: Effects on follicular growth, concentrations of steroids and insulin-like growth factor I, and insulin-like growth factor binding protein activity in follicular fluid of control, lean, and obese gilts. *J Anim Sci* 72:2431–2440, 1994.
42. Gong JG, Bramley T, Webb R. The effect of recombinant bovine somatotropin on ovarian function in heifers: Follicular populations and peripheral hormones. *Biol Reprod* 45:941–949, 1991.
43. Gong JG, Bramley TA, Wilmut I, Webb R. Effect of recombinant bovine somatotropin on the superovulatory response to pregnant mare serum gonadotropin in heifers. *Biol Reprod* 48:1141–1149, 1993.
44. Herrler A, Einspanier R, Schams D, Niemann H. Effect of recombinant bovine somatotropin (rBST) on follicular IGF-I contents and the ovarian response following superovulatory treatment in dairy cows: A preliminary study. *Theriogenology* 41:601–611, 1994.
45. Simpson RB, Chase Jr CC, Spicer LJ, Vernon RK, Hammond AC, Rae DO. Effect of exogenous insulin on plasma and follicular insulin-like growth factor I, insulin-like growth factor binding protein activity, follicular estradiol and progesterone, and follicular growth in superovulated Angus and Brahman cows. *J Reprod Fertil* 102:483–492, 1994.
46. May JV, Bridge AJ, ED Gotcher, Gangrade BK. The regulation of porcine theca cell proliferation in vitro: Synergistic actions of epidermal growth factor and platelet-derived growth factor. *Endocrinology* 131:689–697, 1992.
47. Chakravorty A, Joslyn MI, Davis JS. Characterization of insulin and insulin-like growth factor-I actions in the bovine luteal cell: Regulation of receptor tyrosine kinase activity, phosphatidylinositol-3-kinase, and deoxyribonucleic acid synthesis. *Endocrinology* 133:1331–1340, 1993.
48. Shaw HJ, Boddy SE, Hodges JK. Developmental changes in marmoset granulosa cell responsiveness to insulin-like growth factor-I: Interactions with follicle-stimulating hormone and estradiol. *Biol Reprod* 49:568–576, 1993.
49. Erickson GF, Magoffin DA, Cragun JR, Chang RJ. The effects of insulin and insulin-like growth factors-I

- and -II on estradiol production by granulosa cells of polycystic ovaries. *J Clin Endocrinol Metab* 70: 894–902, 1990.
50. Barreca A, Artini PG, Del Monte P, Ponzani P, Pasquini P, Cariola G, Volpe A, Genazzani AR, Giordano G, Minuto F. In vivo and in vitro effect of growth hormone on estradiol secretion by human granulosa cells. *J Clin Endocrinol Metab* 77:61–67, 1993.
  51. Davoren JB, Kasson BG, Li CH, Hsueh AJW. Specific insulin-like growth factor (IGF) I- and II-binding sites on rat granulosa cells: Relation to IGF action. *Endocrinology* 119:2155–2162, 1986.
  52. Gong JG, McBride D, Bramley TA, Webb R. Effects of recombinant bovine somatotrophin, insulin-like growth factor-I and insulin on bovine granulosa cell steroidogenesis in vitro. *J Endocrinol* 143:157–164, 1994.
  53. Saumande J. Culture of bovine granulosa cells in a chemically defined serum-free medium: The effect of insulin and fibronectin on the response to FSH. *J Steroid Biochem Mol Biol* 38:189–196, 1991.
  54. Meidan R, Girsh E, Blum O, Aberdam E. In vitro differentiation of bovine theca and granulosa cells into small and large luteal-like cells: Morphological and functional characteristics. *Biol Reprod* 43:913–921, 1990.
  55. Spicer LJ, Echternkamp SE, Canning SF, Hammond JM. Relationship between concentrations of immunoreactive insulin-like growth factor-I in follicular fluid and various biochemical markers of differentiation in bovine antral follicles. *Biol Reprod* 39:573–580, 1988.
  56. Spicer LJ, Enright WJ, Murphy MG, Roche JF. Effect of dietary intake on concentrations of insulin-like growth factor-I in plasma and follicular fluid, and ovarian function in heifers. *Domest Anim Endocrinol* 8:431–437, 1991.
  57. Echternkamp SE, Spicer LJ, Gregory KE, Canning SF, Hammond JM. Concentrations of insulin-like growth factor-I in blood and ovarian follicular fluid in cattle selected for twins. *Biol Reprod* 43:8–14, 1990.
  58. Kirby CJ, Armstrong JD, Huff BG, Stanko RL, Harvey RW, Heimer EP, Campbell RM. Changes in serum somatotropin, somatotropin mRNA, and serum and follicular insulin-like growth factor-I in response to feed restriction in cows actively immunized against growth hormone-releasing factor. *J Anim Sci* 71:3033–3042, 1993.
  59. Spicer LJ, Enright WJ. Concentrations of insulin-like growth factor-I and steroids in follicular fluid of preovulatory bovine follicles: Effect of daily injections of a growth hormone-releasing factor analog and/or thyrotropin-releasing hormone. *J Anim Sci* 69:1133–1139, 1991.
  60. Lucy MC, Byatt JC, Curran TL, Curran DF, Collier RJ. Placental lactogen and somatotropin: hormone binding to the corpus luteum and effects on the growth and functions of the ovary in heifers. *Biol Reprod* 50:1136–1144, 1994.
  61. Veldhuis JD, Kolp JA, Toaff ME, Strauss JF III, Demers LM. Mechanisms subserving the trophic actions of insulin on ovarian cells. *J Clin Invest* 72:1046–1057, 1983.
  62. Howard HJ, Ford JJ. Differential steroidogenic response of subpopulations of porcine granulosa cells to insulin-like growth factor-I (IGF-I) or IGF-I analogs. *Biol Reprod* 51:108–115, 1994.
  63. Samaras SE, Guthrie HD, Barber JA, Hammond JM. Expression of the mRNAs for the insulin-like growth factors and their binding proteins during development of porcine ovarian follicles. *Endocrinology* 133: 2395–2398, 1993.
  64. Cox NM, Meurer KA, Carlton CA, Tubbs RC, Mannis DP. Effect of diabetes mellitus during the luteal phase of the oestrous cycle on preovulatory follicular function, ovulation and gonadotrophins in gilts. *J Reprod Fertil* 101:77–86, 1994.
  65. Meurer KA, Cox NM, Matamoros IA, Tubbs RC. Decreased follicular steroids and insulin-like growth factor-I and increased atresia in diabetic gilts during follicular growth stimulated with PMSG. *J Reprod Fertil* 91:187–196, 1991.
  66. Veldhuis JD, Rodgers RJ, Furlanetto RW. Synergistic actions of estradiol and the insulin-like growth factor somatomedin C on swine ovarian (granulosa) cells. *Endocrinology* 119:530–538, 1986.
  67. Veldhuis JD, Rodgers RJ. Mechanisms subserving the steroidogenic synergism between follicle-stimulating hormone and insulin-like growth factor-I (somatomedin C): Alterations in cellular sterol metabolism in swine granulosa cells. *J Biol Chem* 262:7658–7664, 1987.
  68. Schams D, Koll R, Li CH. Insulin-like growth factor-I stimulates oxytocin and progesterone production by bovine granulosa cells in culture. *J Endocrinol* 116:97–100, 1988.
  69. Holtorf A-P, Furuya K, Ivell R, McArdle CA. Oxytocin production and oxytocin messenger ribonucleic acid levels in bovine granulosa cells are regulated by insulin and insulin-like growth factor-I: Dependence on development status of the ovarian follicle. *Endocrinology* 125:2612–2620, 1989.
  70. Garmey JC, Day RN, Day KH, Veldhuis JD. Mechanisms of regulation of ovarian sterol metabolism by

- insulin-like growth factor type II: *In vitro* studies with swine granulosa cells. *Endocrinology* 133:800–808, 1993.
71. McArdle CA, Kohl C, Rieger K, Groner I, Wehrenberg U. Effects of gonadotropins, insulin and insulin-like growth factor I on ovarian oxytocin and progesterone production. *Mol Cell Endocrinol* 78:211–220, 1991.
  72. Li Z, Johnson AL. Regulation of P450 cholesterol side-chain cleavage messenger ribonucleic acid expression and progesterone production in hen granulosa cells. *Biol Reprod* 49:463–469, 1993.
  73. Spicer LJ, Crowe MA, Prendiville DJ, Goulding D, Enright WJ. Systemic but not intraovarian concentrations of insulin-like growth factor-I are affected by short-term fasting. *Biol Reprod* 46:920–925, 1992.
  74. Roberts AJ, Skinner MK. Hormonal regulation of thecal cell function during antral follicle development in bovine ovaries. *Endocrinology* 127:2907–2917, 1990.
  75. Stewart RE, Spicer LJ, Hamilton T, Keefer BE. Effects of insulin-like growth factor-I and insulin on proliferation, and basal and LH-induced steroidogenesis of thecal cells: Involvement of glucose and receptors for IGF-I and LH. *J Anim Sci* (in press).
  76. Caubo B, DeVinna RS, Tonetta SA. Regulation of steroidogenesis in cultured porcine theca cells by growth factors. *Endocrinology* 125:321–326, 1989.
  77. Morley P, Calaresu FR, Barbe GJ, Armstrong DT. Insulin enhances luteinizing hormone-stimulated steroidogenesis by porcine theca cells. *Biol Reprod* 40:735–743, 1989.
  78. Magoffin DA, Weitsman SR. Insulin-like growth factor-I stimulates the expression of 3 $\beta$ -hydroxysteroid dehydrogenase messenger ribonucleic acid in ovarian theca-interstitial cells. *Biol Reprod* 48:1166–1173, 1993.
  79. Magoffin DA, Weitsman SR. Effect of insulin-like factor-I on cholesterol side-chain cleavage cytochrome P450 messenger ribonucleic acid expression in ovarian theca-interstitial cells stimulated to differentiate in vitro. *Mol Cell Endocrinol* 46:45–51, 1993.
  80. Veldhuis JD, Nestler JE, Strauss JF III, Garmey J, Juchter D. The insulin-like growth factor somatomedin-C, modulates low density lipoprotein metabolism by swine granulosa cells. *Endocrinology* 121:340–346, 1987.
  81. Urban RJ, Garmey JC, Shupnik MA, Veldhuis JD. Insulin-like growth factor Type I increases concentrations of messenger ribonucleic acid encoding cytochrome P450 cholesterol side-chain cleavage enzyme in primary cultures of porcine granulosa cells. *Endocrinology* 127:2481–2488, 1990.
  82. Magoffin DA, Kurtz KM, Erickson GF. Insulin-like growth factor selectively stimulates cholesterol side-chain cleavage expression in ovarian theca-interstitial cells. *Mol Endocrinol* 4:489–496, 1990.
  83. McArdle CA, Holtorf AP. Oxytocin and progesterone release from bovine corpus luteal cells in culture: Effects of insulin-like growth factor-I, insulin, and prostaglandins. *Endocrinology* 124:1278–1286, 1989.
  84. Ladenheim RG, Tesone M, Charreau EH. Insulin action and characterization of insulin receptors in rat luteal cells. *Endocrinology* 115:752–756, 1984.
  85. Sauerwein H, Miyamoto A, Gunther J, Meyer HHD, Schams D. Binding and action of insulin-like growth factors and insulin in bovine luteal tissue during the oestrous cycle. *J Reprod Fertil* 96:103–115, 1992.
  86. Constantino CX, Keys PL, Kostyo JL. Insulin-like growth factor-I stimulates steroidogenesis in rabbit luteal cells. *Endocrinology* 128:1702–1708, 1991.
  87. Parmer TG, Roberts CT Jr, LeRoith D, Adashi EY, Khan I, Solan N, Nelson S, Zilberstein M, Gibori G. Expression, action, and steroidal regulation of insulin-like growth factor-I (IGF-I) and IGF-I receptor in the rat corpus luteum: Their differential role in the two cell populations forming the corpus luteum. *Endocrinology* 129:2924–2932, 1991.
  88. Spicer LJ, Tucker WB, Adams GD. Insulin-like growth factor-I in dairy cows: relationships among energy balance, body condition, ovarian activity, and estrous behavior. *J Dairy Sci* 73:929–937, 1990.
  89. Spicer LJ, Vernon RK, Tucker WB, Wettemann RP, Hogue JF, Adams GD. Effects of inert fat on energy balance, plasma concentrations of hormones, and reproduction in dairy cows. *J Dairy Sci* 76:2664–2673, 1993.
  90. Hernandez ER, Resnick CE, Holtzclaw WD, Payne DW, Adashi EY. Insulin as a regulator of androgen biosynthesis by cultured rat ovarian cells: Cellular mechanism(s) underlying physiological and pharmacological hormonal actions. *Endocrinology* 122:2034–2043, 1988.
  91. Hernandez ER, Resnick CE, Svoboda ME, Van Wyk JJ, Payne DW, Adashi EY. Somatomedin-C/insulin-like growth factor I as an enhancer of androgen biosynthesis by cultured rat ovarian cells. *Endocrinology* 122:1603–1612, 1988.
  92. Simone DA, Mahesh VB. An autoregulatory process for androgen production in rat thecal-interstitial cells. *Biol Reprod* 48:46–46, 1993.
  93. Bergh C, Carlsson B, Olsson J, Selleskog U, Hillensjo T. Regulation of androgen production in cultured human thecal cells by insulin-like growth factor I and insulin. *Fertil Steril* 59:323–331, 1993.

94. Hammond JM, Veldhuis JD, Seale TW, Rechler MM. Intraovarian regulation of granulosa-cell replication. In: *Intraovarian Control Mechanisms*, Channing CP, Segal SJ (eds), Plenum Press, New York, pp. 341–356, 1982.
95. Hammond JM, Baranao JLS, Skaleris D, Knight AB, Romanus JA, Rechler MM. Production of insulin-like growth factors by ovarian granulosa cells. *Endocrinology* 117:2553–2555, 1985.
96. Jesionowska H, Hemmings R, Guyda HJ, Posner BI. Determination of insulin and insulin-like growth factors in the ovarian circulation. *Fertil Steril* 53:88–91, 1990.
97. Spicer LJ, Tucker KE, Henderson KA, Duby RT. Concentrations of insulin-like growth factor-I in follicular fluid and blood plasma of mares during early and late oestrus. *Anim Reprod Sci* 25:57–65, 1991.
98. Spicer LJ, Echternkamp SE, Wong EA, Hamilton DT, Vernon RK. Serum hormones, follicular fluid steroids, insulin-like growth factors and their binding proteins, and ovarian IGF mRNA in sheep with different ovulation rates. *J Anim Sci* 73:1152–1163, 1995.
99. Hammond JM, Hsu C-J, Klindt J, Ainsworth L, Downey BR, Tsang BK. Gonadotropins increase concentrations of immunoreactive insulin-like growth factor-I in porcine follicular fluid in vivo. *Biol Reprod* 38:304–308, 1988.
100. Samaras SE, Hagen DR, Bryan KA, Mondschein JS, Canning SF, Hammond JM. Effects of growth hormone and gonadotropin on the insulin-like growth factor system in the porcine ovary. *Biol Reprod* 50:178–186, 1994.
101. Diamond MP, Webster BW, Carr RK, Wentz AC, Osteen KG. Human follicular fluid insulin concentrations. *J Clin Endocrinol Metab* 61:990–992, 1984.
102. Murphy LJ, Bell GI, Friesen HG. Tissue distribution of insulin-like growth factor-I and II messenger ribonucleic acid in the adult rat. *Endocrinology* 120:1279–1282, 1987.
103. Voutilainen R, Miller WL. Coordinate tropic hormone regulation of mRNAs for insulin-like growth factor II and the cholesterol side-chain-cleavage enzyme, P450 scc in human steroidogenic tissues. *Proc Natl Acad Sci USA* 84:1590–1594, 1987.
104. Geisthovel F, Moretti-Rojas I, Asch RH, Rojas FJ. Expression of insulin-like growth factor-II (IGF-II) messenger ribonucleic acid (mRNA), but not IGF-I mRNA, in human preovulatory granulosa cells. *Hum Reprod* 4:899–902, 1989.
105. Hernandez ER, Roberts CT, Jr, LeRoith D, Adashi EY. Rat ovarian insulin-like growth factor-I (IGF-I) gene expression is granulosa cell-selective: 5'-Untranslated mRNA variant representation and hormonal regulation. *Endocrinology* 125:572–574, 1989.
106. Oliver JE, Aitman TJ, Powell JF, Wilson CA, Clayton RN. Insulin-like growth factor I gene expression in the rat ovary is confined to the granulosa cells of developing follicles. *Endocrinology* 124:2671–2679, 1989.
107. Einspanier R, Miyamoto A, Schams D, Muller M, Brem G. Tissue concentration, mRNA expression and stimulation of IGF-I in luteal tissue during the oestrous cycle and pregnancy of cows. *J Reprod Fertil* 90:439–445, 1990.
108. Charlton ST, Cameron BJ, Glimm DR, Foxcroft GR, Kennelly JJ. Insulin-like growth factor (IGF-I) gene expression in porcine ovarian tissue. *Can J Anim Sci* 73:253–257, 1993.
109. Ramasharma K, Li CH. Human pituitary and placental hormones control human insulin-like growth factor II secretion in human granulosa cells. *Proc Natl Acad Sci USA* 84:2643–2647, 1987.
110. El-Roeiy A, Chen X, Roberts VJ, LeRoith D, Roberts CT, Jr, Yen SSC. Expression of insulin-like growth factor-I (IGF-I) and IGF-II and the IGF-I, IGF-II, and insulin receptor genes and localization of the gene products in the human ovary. *J Clin Endocrinol Metab* 77:1411–1418, 1993.
111. El-Roeiy A, Chen X, Roberts VJ, Shimasaki S, Ling N, LeRoith D, Roberts CT Jr, Yen SSC. Expression of the genes encoding the insulin-like growth factors (IGF-I and II), the IGF and insulin receptors, and IGF-binding proteins-1-6 and the localization of their gene products in normal and polycystic ovary syndrome ovaries. *J Clin Endocrinol Metab* 78:1488–1496, 1994.
112. Yoshimura Y, Iwashita M, Karube M, Oda T, Akiba M, Shiokawa S, Ando M, Yoshinaga A, Nakamura Y. Growth hormone stimulates follicular development by stimulating ovarian production of insulin-like growth factor-I. *Endocrinology* 135:887–894, 1994.
113. Gates GS, Bayer S, Seibel M, Poretsky L, Flier JS, Moses AC. Characterization of insulin-like growth factor binding to human granulosa cells obtained during in vitro fertilization. *J Recept Res* 7:855–902, 1987.
114. Samoto T, Maruo T, Matsuo H, Katayama K, Barnea ER, Mochizuki M. Altered expression of insulin and insulin-like growth factor-I receptors in follicular and stromal compartments of polycystic ovaries. *Endocrine J* 40:413–424, 1993.
115. Samoto T, Mauro T, Ladines-Llave CA, Matsuo H, Deguchi J, Barnea ER, Mochizuki M. Insulin receptor expression in follicular and stromal compartments of the human ovary over the course of follicular growth, regression and atresia. *Endocrine J* 40:715–726, 1993.

116. Talavera, F. and Menon K.M.J. Studies on rat luteal cell response to insulin-like growth factor I (IGF-I): Identification of specific cell membrane receptor for IGF-I in the luteinized rat ovary. *Endocrinology* 129:1340–1346, 1991.
117. Jarrett JC, Jr, Ballejo G, Tsisbris JCM, Spellacy WN. Insulin binding to human ovaries. *J Clin Endocrinol Metab* 60:460–463, 1985.
118. Poretsky L, Bhargava G, Kalin MF, Wolf SA. Regulation of insulin receptors in the human ovary: In vitro studies. *J Clin Endocrinol Metab* 67:774–778, 1988.
119. Hernandez ER, Hurwitz A, Vera A, Pellicer A, Adashi EY, LeRoith D, Roberts Jr CT. Expression of the genes encoding the insulin-like growth factors and their receptors in the human ovary. *J Clin Endocrinol Metab* 74:419–425, 1992.
120. Hylka VW, Caubo B, Tonetta SA. Analysis of binding sites for IGF-I on membranes from granulosa cells of small, medium, and large porcine follicles. In: *Growth Factors and the Ovary*, Hishfield AN (ed). Plenum Press, New York, p. 163–168, 1989.
121. Wandji S-A, Pelletier G, Sirard M-A Ontogeny and cellular localization of  $^{125}$ I-labeled insulin-like growth factor,  $^{125}$ I-labeled follicle-stimulating hormone and  $^{125}$ I-labeled human chorionic gonadotropin binding activities in ovaries from bovine fetuses and neonatal calves. *Biol Reprod* 47:814–822, 1992.
122. Zhou J, Chin E, Bondy C. Cellular pattern of insulin-like growth factor-I (IGF-I) and IGF-I receptor gene expression in the developing and mature ovarian follicle. *Endocrinology* 129:3281–3288, 1991.
123. Metcalf MG. Estimation of viability of bovine granulosa cells. *J Reprod Fertil* 65:425–429, 1982.
124. Buck PA, Schomberg DW. A serum-free defined culture system which maintains follicle-stimulating hormone responsiveness and differentiation of porcine granulosa cells. *Biol Reprod* 36:167–174, 1987.
125. Monget P, Monniaux D, Durand P. Localization, characterization, and quantification of insulin-like growth factor-I binding sites in the ewe ovary. *Endocrinology* 125:2486–2493, 1989.
126. Zhou J, Bondy C. Anatomy of the human ovarian insulin-like growth factor system. *Biol Reprod* 48:467–482, 1993.
127. Balboni GC, Vannelli GB, Barni T, Orlando C, Serio M. Transferrin and somatomedin C receptors in the human ovarian follicles. *Fertil Steril* 48:796–801, 1987.
128. Urban RJ, Bodenburg YH, Nagamani M, Peirce J. Dexamethasone potentiates insulin-like growth factor-I actions in porcine granulosa cells. *Am J Physiol* 267:E115–E123, 1994.
129. Aashi EY, Resnick CE, Hernandez ER, Svoboda ME, Van Wyk JJ. Follicle-stimulating hormone enhances somatomedin C binding to cultured rat granulosa cells (evidence for cAMP dependence). *J Biol Chem* 261:3923–3926, 1986.
130. Adashi EY, Resnick CE, Hernandez ER, Svoboda ME, Van Wyk JJ. Characterization and regulation of a specific cell membrane receptor for somatomedin-C/insulin-like growth factor I in cultured rat granulosa cells. *Endocrinology* 122:194–201, 1988.
131. Teissier M-P, Monget P, Monniaux D, Durand P. Changes in insulin-like growth factor-II/mannose-6-phosphate receptor during growth and atresia of ovine ovarian follicles. *Biol Reprod* 50:111–119, 1994.
132. Clemmons DR. Insulin-like growth factor binding proteins. In: *Insulin-Like Growth Factors: Molecular and Cellular Aspects*, LeRoith D (ed). CRC Press, Boca Raton, FL, pp. 151–179, 1991.
133. Clemmons DR. IGF binding proteins and their functions. *Mol Reprod Devel* 5:368–375, 1993.
134. Rechler MM. Insulin-like growth factor binding proteins. *Vitam Horm* 47:1–114, 1993.
135. Margot JB, Binkert C, Mary JL, Landwehr J, Heinrich G, Schwander J. A low molecular weight insulin-like growth factor binding protein from the rat: cDNA cloning and tissue distribution of its messenger RNA. *Mol Endocrinol* 3:1053–1060, 1989.
136. Shimasaki S, Koba A, Mercado M, Shimonaka M, Ling N. Complementary DNA structure of a high molecular weight rat insulin-like growth factor binding protein (IGF-BP-3) and tissue distribution of its mRNA. *Biochem Biophys Res Commun* 165:907–912, 1989.
137. Ui M, Shimonaka M, Shimasaki S, Ling N. An insulin-like growth factor-binding protein in ovarian follicular fluid blocks follicle-stimulating hormone-stimulated steroid production by ovarian granulosa cells. *Endocrinology* 125:912–916, 1989.
138. Bicsak TA, Ling N, DePaolo LV. Ovarian intrabursal administration of insulin-like growth factor-binding protein inhibits follicle rupture in gonadotropin-treated immature rats. *Biol Reprod* 44:599–603, 1991.
139. Monget P, Monniaux D, Pisselet C, Durand P. Changes in insulin-like growth factor-I (IGF-I), IGF-II, and their binding proteins during growth and atresia of ovine ovarian follicles. *Endocrinology* 132:1438–1446, 1993.
140. Mason HD, Willis D, Holly JMP, Cwyfan-Hughes SC, Seppala M, Franks S. Inhibitory effects of insulin-like growth factor-binding proteins on steroidogenesis by human granulosa cells in culture. *Mol Cell Endocrinol* 89:R1–R4, 1992.
141. Liu X, Malkowski M, Guo Y, Erickson GF, Shimasaki S, Ling N. Development of specific antibodies to

- rat insulin-like growth factor-binding proteins (IGFBP-2 to -6): Analysis of IGFBP production by rat granulosa cells. *Endocrinology* 132:1176–1183, 1993.
142. Echternkamp SE, Howard HJ, Roberts AJ, Grizzle J, Wise T. Relationship among concentrations of steroids, insulin-like growth factor-I, and insulin-like growth factor binding proteins in ovarian follicular fluid of beef cattle. *Biol Reprod* 51:971–981, 1994.
  143. Mondschein JS, Etherton TD, Hammond JM. Characterization of insulin-like growth factor-binding proteins of porcine ovarian follicular fluid. *Biol Reprod* 44:315–320, 1991.
  144. Howard HJ, Ford JJ. Relationships among concentrations of steroids, inhibin, insulin-like growth factor (IGF-I), and IGF-binding proteins during follicular development in weaned sows. *Biol Reprod* 47:193–201, 1992.
  145. San Roman GA, Magoffin DA. Insulin-like growth factor-binding proteins in healthy and atretic follicles during natural menstrual cycles. *J Clin Endocrinol Metab* 76:625–632, 1992.
  146. Fielder PJ, Pham H, Adashi EY, Rosenfeld RG. Insulin-like growth factors (IGFs) block FSH-induced proteolysis of IGF-binding protein-5 (BP-5) in cultured rat granulosa cells. *Endocrinology* 133:415–418, 1993.
  147. Grimes RW, Hammond JM. Proteolytic degradation of insulin-like growth factor (IGF)-binding protein-3 by porcine ovarian granulosa cells in culture: Regulation by IGF-I. *Endocrinology* 134:337–343, 1994.
  148. Grimes RW, Samaras SE, Hammond JM. Divergent actions of prostaglandins-E<sub>2</sub> and -F<sub>2</sub>  $\alpha$  on the regulation of insulin-like growth factor-binding protein-3 in luteinized granulosa cells. *Endocrinology* 132:1414–1416, 1993.
  149. Samaras SE, Hagen DR, Shimasaki S, Ling N, Hammond JM. Expression of insulin-like growth factor binding protein (IGFBP)-2 and -3 mRNA in the porcine ovary: localization and physiological changes. *Endocrinology* 130:2739–2744, 1992.
  150. Grimes RW, Barber JA, Shimasaki S, Ling N, Hammond JM. Porcine ovarian granulosa cells secrete insulin-like growth factor-binding proteins-4 and -5 and express their messenger ribonucleic acids: Regulation by follicle-stimulating hormone and insulin-like growth factor-I. *Biol Reprod* 50:695–701, 1994.
  151. Nakatani A, Shimasaki S, Erickson GF, Ling N. Tissue-specific expression of four insulin-like growth factor-binding proteins (1, 2, 3, and 4) in the rat ovary. *Endocrinology* 129:1521–1529, 1991.
  152. Ricciarelli E, Hernandez ER, Hurwitz A, Kokia E, Rosenfeld RJ, Schwander J, Adashi EY. The ovarian expression of the antagonodotropic insulin-like growth factor binding protein-2 is theca-interstitial cell-selective: Evidence for hormonal regulation. *Endocrinology* 129:2266–2268, 1991.
  153. Erickson GF, Nakatani A, Ling N, Shimasaki S. Cyclic changes in insulin-like growth factor-binding protein-4 messenger ribonucleic acid in the rat ovary. *Endocrinology* 130:625–636, 1992.
  154. Erickson GF, Nakatani A, Ling N, Shimasaki S. Localization of insulin-like growth factor-binding protein-5 messenger ribonucleic acid in rat ovaries during the estrous cycle. *Endocrinology* 130:1867–1878, 1992.
  155. Ricciarelli E, Hernandez ER, Tedeschi C, Botero LF, Kokia E, Rohan RM, Rosenfeld RG, Albiston AL, Herington AC, Adashi EY. Rat ovarian insulin-like growth factor binding protein-3: A growth hormone-dependent theca-interstitial cell-derived antagonodotropin. *Endocrinology* 130:3092–3094, 1992.
  156. Erickson GF, Nakatani A, Ling N, Shimasaki S. Insulin-like growth factor binding protein-3 gene expression is restricted to involuting corpora lutea in rat ovaries. *Endocrinology* 133:1147–1157, 1993.
  157. Jalkanen J, Suikkari A, Koistinen R, Butzow R, Ritvos O, Seppala M, Ranta T. Regulation of insulin-like growth factor-binding protein-1 production in human granulosa-luteal cells. *J Clin Endocrinol Metab* 69:1174–1179, 1989.
  158. Stanko RL, Cohick WS, Shaw DW, Harvey RW, Clemmons DR, Whitacre MD, Armstrong JD. Effect of somatotropin and/or equine chorionic gonadotropin on serum and follicular insulin-like growth factor-I and insulin-like growth factor binding proteins in cattle. *Biol Reprod* 50:290–300, 1994.
  159. Giudice LC, Farrell EM, Pham H, Rosenfeld RG. Identification of insulin-like growth factor-binding protein-3 (IGFBP-3) and IGFBP-2 in human follicular fluid. *J Clin Endocrinol Metab* 71:1330–1338, 1990.
  160. San Roman GA, Magoffin DA. Insulin-like growth factor binding proteins in ovarian follicles from women with polycystic ovarian disease: Cellular source and levels in follicular fluid. *J Clin Endocrinol Metab* 75:1010–1016, 1992.
  161. Clemmons DR, Underwood LE. Nutritional regulation of IGF-I and IGF binding proteins. *Annu Rev Nutr* 11:393–412, 1991.
  162. Thissen J-P, Ketelslegers J-M, Underwood LE. Nutritional regulation of the insulin-like growth factors. *Endocr Rev* 15:80–101, 1994.
  163. Cataldo NA, Woodruff WD, Giudice LC. Regulation of insulin-like growth factor binding protein pro-

- duction by human luteinizing granulosa cells cultured in defined medium. *J Clin Endocrinol Metab* 76:207–215, 1993.
164. Adashi EY, Resnick CE, Hurwitz A, Ricciarelli E, Hernandez ER, Rosenfeld RG. Ovarian granulosa cell-derived insulin-like growth factor binding proteins: Modulatory role of follicle-stimulating hormone. *Endocrinology* 128:754–760, 1991.
  165. Adashi EY, Resnick CE, Hernandez ER, Hurwitz A, Rosenfeld RG. Follicle-stimulating hormone inhibits the constitutive release of insulin-like growth factor binding proteins by cultured rat ovarian granulosa cells. *Endocrinology* 126:1305–1307, 1990.
  166. Adashi EY, Resnick CE, Tedeschi C, Rosenfeld RG. A kinase-mediated regulation of granulosa cell-derived insulin-like growth factor binding proteins (IGFBPs): Disparate response sensitivities of distinct IGFBP species. *Endocrinology* 132:1463–1468, 1993.
  167. Dor J, Costritsci N, Pariente C, Rabinovic J, Mashlach S, Lunenfeld B, Kaneti H, Seppala M, Koistinen R, Karasik A. Insulin-like growth factor-I and follicle-stimulating hormone suppress insulin-like growth factor binding protein-I secretion by human granulosa-luteal cells. *J Clin Endocrinol Metab* 75:969–971, 1992.
  168. Rohan RM, Ricciarelli E, Kiefer MC, Resnick CE, Adashi EY. Rat ovarian insulin-like growth factor-binding protein-6: A hormonally regulated theca-interstitial-selective species with limited antigonadotropic activity. *Endocrinology* 132:2507–2512, 1993.
  169. Spicer LJ, Hanrahan JP, Zavy MT, Enright WJ. Relationship between ovulation rate and concentrations of insulin-like growth factor-I in plasma during the oestrous cycle in various genotypes of sheep. *J Reprod Fertil* 97:403–409, 1993.
  170. Leeuwenberg BR, Hurst P, McNatty KP. Expression of insulin-like growth factor-I (IGF-I) in the sheep ovary. *Biol Reprod* 48[Suppl 1]:184, 1993.
  171. Murphy MG, Enright WJ, Crowe MA, McConnell K, Spicer LJ, Boland MP, Roche JF. Effect of dietary intake on pattern of growth of dominant follicles during the oestrous cycle in beef heifers. *J Reprod Fertil* 92:333–338, 1991.
  172. Simmen FA, Simmen RCM. Peptide growth factors and proto-oncogenes in mammalian conceptus development. *Biol. Reprod.* 44:1–5, 1991.
  173. Simmen RCM, Simmen FA, Hofig A, Farmer SJ, Bazer FW. Hormonal regulation of insulin-like growth factor gene expression in pig uterus. *Endocrinology* 127:2166–2174, 1990.
  174. Geisert RD, Lee, CY, Simmen FA, Zavy MT, Fliss AE, Bazer FW, Simmen RCM. Expression of messenger RNAs encoding insulin-like growth factor-I, -II and insulin-like growth factor binding protein-2 in bovine endometrium during the estrous cycle and early pregnancy. *Biol Reprod* 45:975–983, 1991.
  175. Stevenson KR, Gilmour RS, Wathes DC. Localization of insulin-like growth factor-I (IGF-I) and -II messenger ribonucleic acid and type I IGF receptors in the ovine uterus during the estrous cycle and early pregnancy. *Endocrinology* 134:1655–1664, 1994.
  176. Murphy LH, Freisen HG. Differential effects of estrogen and growth hormone on uterine hepatic insulin-like growth factor I gene expression in the ovariectomized hypophysectomized rat. *Endocrinology* 122:325–332, 1988.
  177. Murphy LJ. Estrogen induction of insulin-like growth factors and myc proto-oncogene expression in the uterus. *J Steroid Biochem Molec Biol* 40:223–230, 1991.
  178. Rieger D, Walton JS, Goodwin ML, Johnson WH. The effect of co-treatment with recombinant bovine somatotropin on plasma progesterone concentration and number of embryos collected from superovulated Holstein heifers. *Theriogenology* 35:863–867, 1991.
  179. Cogie Y, Poulin N, Guerin Y, Martinat N. Administration of exogenous hormone in early follicular phase enhance embryo production in sheep. *Proc 12th Int Congr Anim Reprod* 2:785–787, 1992.
  180. Bryan KA, Hammond JM, Canning S, Mondschein J, Carbaugh DE, Clark AM, Hagen DR. Reproductive and growth responses of gilts to exogenous porcine pituitary growth hormone. *J Anim Sci* 67:196–205, 1989.
  181. Gallo GF, Block E. Effect of recombinant bovine somatotropin on hypophyseal and ovarian functions of lactating dairy cows. *Can J Anim Sci* 71:343–353, 1991.
  182. Fotsis T, Murphy C, Gannon F. Nucleotide sequence of the bovine insulin-like growth factor I (IGF-I) and its IGF-IA precursor. *Nucleic Acids Res* 18:676, 1990.